

Calcareous nannoplankton from the Seoguipo Formation of Cheju Island, Korea and its paleoceanographic implications

SONGSUK YI¹, HYESU YUN² and SUN YOON³

¹Technical Department, Korea National Oil Corporation, 1588-14, Kwanyang-dong, Dongan-gu, Anyang, Kyungki-do 431-711, Korea

²Department of Geology, College of Natural Sciences, Chungnam National University, Taejeon 305-764, Korea

³Department of Geology, College of Natural Sciences, Pusan National University, Pusan 609-735, Korea

Received 25 June 1998 ; Revised manuscript accepted 2 November 1998

Abstract. Twenty species of calcareous nannofossils belonging to 11 genera are identified from the Seoguipo Formation in Cheju Island, Korea. On the basis of the marker species, the Seoguipo Formation is biostratigraphically assigned to the *Pseudoemiliana lacunosa* Zone (NN19), which corresponds to the combined zones of *Emiliana annula*–*Emiliana ovata* (CN13a–CN14a) of the latest Pliocene and Early Pleistocene. Generally, cold-water species are dominant in the lower part, and warm-water ones in the upper part. This suggests that the paleoceanographic condition of the study area changed from a cooling to a warm phase. The change in floral composition and abundance of specific species allows recognition of four ecostratigraphic zones in the Seoguipo Formation and the migration of an oceanographic frontal boundary. According to nannofossil distribution in the study area, the position of an oceanographic boundary between warmer water and cooler water appeared to have oscillated north–south over the Korea Strait and Cheju Island in response to glacial and interglacial cycles. The geologic time of the interpreted paleoceanographical changes determined by nannofossil biochronology agrees well with the results obtained from the Japan Sea (East Sea) and Japan–Sea side of Japan.

Key words: Bio- and ecostratigraphy, Cheju Island, Korea, nannoplankton, paleoceanography, Seoguipo Formation.

Introduction

Cheju Island, located 90 km off the southern coast of Korea at 33°12'–33°34'N, 126°10'–126°58'E, was formed by Late Cenozoic volcanic activity (Figure 1). The Seoguipo Formation, a highly fossiliferous marine formation, seems to have been deposited just before the onset of Quaternary volcanism, since it underlies Pleistocene volcanic rocks that cover most of the island (Table 1). The Seoguipo Formation is mainly composed of sandstone with a minor amount of interbedded conglomerate and mudstone. It is well developed at the subsurface level all over the island and is exposed only on the southern coast of the island at Seoguipo. The subsurface Seoguipo Formation is encountered in core samples of numerous bore holes drilled in the island.

Yokoyama (1923) first reported molluscan fossils from these sedimentary deposits, designating their geologic age as Upper Pliocene. Haraguchi (1931), who described diverse invertebrate faunas such as mollusks, brachiopods,

echinoids, corals and fish teeth, named this fossiliferous sedimentary formation the Seikiho (Seoguipo) Formation of the Pleistocene epoch. After Haraguchi's research, numerous works including detailed field survey, paleontological, magnetostratigraphic, and sedimentological studies were carried out for determination of the geologic age, depositional environment, and stratigraphical zonation (Kim, 1972; Won, 1975; Lee, M.W., 1982a, 1982b; Kim, 1984; Nomura, 1984; Yun *et al.*, 1987; Min *et al.*, 1986; You *et al.*, 1987; Yoon, 1988; Tamanyu, 1990; Lee E.H., 1990; Yoon *et al.*, 1995; Kang, 1995). However, no consensus was achieved for the geologic age of the Seoguipo Formation which was assigned to different ages from Pliocene to Pleistocene (Table 1). Several studies analyzed paleoclimate and paleoceanographic conditions of the Seoguipo Formation based on molluscan fauna and isotope data (Yoon, 1988; Amano, 1994; Park *et al.*, 1994; Woo *et al.*, 1995). The results for the water temperature are also contradictory, reporting warm or cold glacial environment during deposition of the formation.




Figure 1. Geologic map of Cheju Island with sample locality (after Yoon *et al.*, 1995).

Cheju Island is located in the oceanic pathway to the Japan Sea. This area was paleoclimatologically and tectonically sensitive and often affected by small- or large-scale changes in the tectonic (Inoue, 1982) or climatological settings. The climatic or sea-level changes may cause subsidence or uplift which in turn affects the system of current and the biofacies. Therefore, a biofacies analysis may deductively provide a clue to the climatic and tectonic history of this area, which is not only important for the development of Cheju Island and the Korea Strait, but also for the evolution of the Japan Sea during the late Pliocene and Pleistocene (Muza, 1992 ; Rahman, 1992).

Analysis of calcareous nannofossil assemblage is the best means to resolve the debateable claims made concerning the geologic age and environment of the Seoguiipo Formation, since it may enable fine zonation and reconstruction of the paleoenvironmental and geologic events. Therefore, our study primarily aims to establish a biozonation and ecostratigraphy based on nannoplankton, and to investigate the frequency and nature of paleoceanographic changes during deposition of the Seoguiipo Formation. The results may help in interpretation of the paleoclimatic and tectonic history of Cheju Island and the neighboring area.

Geologic setting of the Seoguiipo Formation

The Seoguiipo Formation is divided into 7 distinctive lithologic units in ascending order (Lee, 1990 ; Figure 2). Litho-unit I of about 3.5 m thickness is composed of laminated yellowish gray siltstones in the lower part, and massive fine-grained gray sandstones with fossil shells and volcanic clasts in the upper part. Litho-unit II overlying the Litho-unit I with a diastem is about 5 m thick, and consists of alternating brownish medium-grained, parallel-stratified sandstones and thin-laminated mudstones. The upper part of this unit with a thickness of about 2 m is highly bioturbated and contains a few articulated bivalve shells. Litho-unit III with a thickness of about 15 m is relatively homogeneous. It is composed of poorly consolidated massive sandstones. The lower boundary of this unit nearly coincides with the Pliocene-Pleistocene boundary. Except for the middle part this unit is highly fossiliferous and bioturbated. Litho-unit IV with a thickness of about 2 m is composed of massive yellowish-gray fine-grained sandstones and highly bioturbated mudstones. Litho-unit V, which is about 10 m in thickness, overlies Litho-unit IV with a diastem. It is lithologically very variable showing cross-laminated shell conglomerate layers, medium- to coarse-grained sandstones intercalated with mudstones, and massive, poorly sorted gray

Table 1. Stratigraphic classifications of volcanics and sedimentary strata in Cheju Island (The sources of the radiometric ages are indicated in brackets : a. Ahn *et al.*, 1995 ; b. Yoon *et al.*, 1995 ; c. Lee, 1994 ; d. Won *et al.*, 1986 ; e. Sameshima *et al.*, 1988).

Age	Haraoguchi, K. (1931)	Won, C.K. (1975)	Lee, M.W. (1982)	Yun, S.K. <i>et al.</i> (1987)	Tamaryu, S. (1990)	Yoon, S. <i>et al.</i> (1995)
Holocene	Flood Deposits Shell Sand Bed Wacke Bed	Shell sand Formation Parasitic volcanic ejectas	1007, 1002 activities Groups of small basalt cones	1007, 1002 activities Dongnam palaeosol	1002, 1007 activities Shell and Formation Scoria vol. cones	Sinyangni Formation [4780 ± 60 y.B.P.] ^e
	Gunzan Basalt Sukido Basalt Kanasan Basalt Aphanitic Basalt Augite Basalt Feldspar Basalt Saishu Basalt Alkali-basalt Trachy-andesite	Baegnogdam Basalt Hallasan Trachy-andesite Hallasan Basalt Seongpanak Basalt Shungri Basalt Beobjeongni Trachyte Hahyori Basalt Jeju Basalt	Sinyangni Formation Hallasan Trachyte Hallasan hawaite Seonpanak hawaite Shungri hawaite Beobjeongni mugearite Hahyori hawaite Jeju hawaite	Dongnam palaeosol Basalt flows along the flanks of Hallasan [0.07 ± 0.04 – 0.035 ± 0.014 Ma] Sinyangni Formation Hallasan trachyte Tuff-rings (Songaksan, Ichulbong, Suwolbong) Seogwipo trachyandesite [0.41 ± 0.01 Ma] Tuff-rings (Dansen, Dusan) Seogwipo Formation Pyoseonni basalt [0.63 ± 0.03 Ma] Tuff-rings (Hwasoon, Dansen)	Paeknoktam Hawaite [0.47 ± 0.07 Ma] Hallasan Trachyte [0.07 ± 0.01 Ma] Pyoseonni Alkali Basalt [0.31 ± 0.04 Ma] Hallasan Hawaite [0.52 ± 0.03 Ma] Sogwipo o Hawaite [0.55 ± 0.04 Ma]	Basognogdam Trachyte Group Yeongsil Trachyte Hallasan Trachybasalt Seonpanak Trachybasalt Cheju Trachybasalt
Pleistocene	Homblende-bearing Trachy-andesite Gyojesigaku Lava Kakushugan Lava Sanbosan Lava	Sinyangni Formation Jungmun Trachyte Seongsampo Fm. Hwasun Fm. Seogwipo Trachyte Pyoseonni Basalt	Homblende mugearite Sanbansan Trachyte Jungmun hawaite Seongsampo Formation Seogwipo hawaite Pyoseonni alkali basalt	Pyoseonni basalt [0.63 ± 0.03 Ma] Sanbansan Trachyte [0.67 ± 0.13 Ma] marine sediments basalt (olivine augite basalt) [1.20 ± 0.05 Ma]	Pyoseonni Basalt Group Jungmun Trachybasalt Group Sanbansan Trachyte Seogwipo Formation U Formation (uncemented sediment beds) Gaumdeog Formation Tanna Conglomerate Sanghyo Trachyte Donnaeoko Basalt Pampo Basalt [2.22 ± 0.16 Ma] ^b	Pyoseonni Basalt Group Jungmun Trachybasalt Group Sanbansan Trachyte Group Seogwipo Formation U Formation (uncemented sediment beds) Gaumdeog Formation Tanna Conglomerate Sanghyo Trachyte Donnaeoko Basalt Pampo Basalt [2.22 ± 0.16 Ma] ^b
Early	Saikho Formation Shinto Lava Johangaku Lava Gyojesigaku Lava Kanasan Lava Granite Blocks bearing Volcanic Detritus Bed	Seogwipo Formation Basal basalt	Seogwipo Formation Basal Basalt	Granite in the northern part Tuff in the eastern and southern part	Granite ?	Granite [5814 ± 1.4 Ma] ^a Volcanic sandstones and mudstones Welded tuffs Lapilli tuffs
	Granite ?		Granite ?			




Figure 2. Biozonation and ecostratigraphic zones of the outcrop section of the Seoguipo Formation based on nannofossils.

medium to coarse sandstones. This unit is characterized by a variety of sedimentary structures such as ripple drift laminae, wavy bedding, flaser bedding, graded bedding, and convolute bedding. Litho-unit VI with a thickness of about 14 m is composed of semi-consolidated massive, light-gray siltstones and volcanic ash. Litho-unit VII of about 15 m thickness consists of volcanic conglomerate without any fossil remains. This unit is covered by a basalt flow.

Methods

Twenty-two samples were collected from the outcrop of the Seoguipo Formation (Figure 2, Table 2). Smear slides were prepared following the centrifuging method of Perch-

Nielsen (1985, p. 330-331). Samples were crushed and centrifuged for 30 seconds at 2,000 r.p.m. Slides were examined with a light microscope at $\times 1,250$ magnification and a scanning electron microscope. Preservation and abundance of nannofossils in each sample are denoted as follow: Abundant (A), 1-10 specimens/one field of view; Common (C), 1 specimen/2-10 fields of view; Few (F), 1 specimen/11-100 fields of view; Rare (R), 1 specimen/101-1,000 fields of view; Barren (B), no fossils observed; Good (G): specimens show little effects of dissolution or overgrowth; Moderate (M), specimens show some effects of dissolution or overgrowth.

Table 2. Stratigraphic occurrence of calcareous nannofossil from the Seoguipo Formation.

Pliocene	Pleistocene		Geologic Age																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																								
	Pseudoemiliania lacunosa Zone (NN19)																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																										
CN13a	CN13b	CN14a	Nannofossil zone																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																								
				Ecostratigraphic Zone																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
I	II	III	IV	Formation																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
				Seoguipo																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
85S02	85S03	SGF 1	SGF 2	SGF 3	SGF 4	C	M	B	A	G	C	M	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C	F	R	C

Results


1. Calcareous nannofossil flora

Twenty species of calcareous nannofossils belonging to 11 genera were identified from 22 samples from the Seoguipo Formation (Table 2). Nannofossils were generally common to abundant. However, some species changed their relative abundance throughout the formation. The dominant species are small *Gephyrocapsa* spp. (<3 μ m; Figures 3-1-3, 4-1, 2), *Pseudoemiliania lacunosa* (Figures 3-12, 13, 4-4, 5), *Coccolithus pelagicus* (Figures 3-28, 29, 4-15), small *Reticulofenestra* spp. (Figures 3-8-10, 4-3), *Gephyrocapsa caribbeanica* (3.5–4 μ m; Figures 3-4, 5, 4-7, 8) and *G. oceanica* (>4 μ m; Figures 3-6, 7, 4-9, 10) in ascending order in the formation. The minor species include *Braarudosphaera bigelowii* (Figure 3-32), *Calcidiscus leptoporus* (Figures 3-20a, b, 27, 4-12-14), *C. macintyreii* (Figures 3-25a, b, 26), *Ceratolithus cristatus* (Figures 3-33a, b), *Coccolithus* sp. (Figures 3-19, 23, 24, 4-11), *Helicosphaera carteri* (Figures

3-21, 22, 4-16), *Helicosphaera* sp. (Figures 3-16, 17), *Pontosphaera japonica* (Figures 3-30a, b, 31, 4-17), *P. discopora*, *Reticulofenestra asanoi* (Figure 3-11), *Syracosphaera pulchra* (Figures 3-15, 4-18), *Syracosphaera* sp. (Figure 3-14), *Umbilicosphaera hultburtiana* and *U. sibogae* (Figures 3-18, 4-6).

The nannoflora in the formation shows a transition from cold subpolar to warm subtropical assemblage. Abundance of the typical cold-water indicator *C. pelagicus* varies greatly within the formation. *C. pelagicus* is extremely abundant in the lower part of the formation and gradually decreases upward, becoming few to barren. Subtropical nannofossils such as *G. oceanica* (>4 μ m), *C. leptoporus*, *U. sibogae*, *S. pulchra*, and *C. cristatus* are common in the upper part of the formation.

The assemblage characteristics change in response to the paleoceanographic conditions during deposition of the formation and hence allow recognition of four ecostratigraphic zones in the outcrop section of the Seoguipo Formation (Figure 2). Zone I (SGF 1-2) contains dominant small *Ge-*



phycocapsa spp. ($<3\ \mu\text{m}$), common *P. lacunosa* and small *Reticulofenestra* spp., and few *C. pelagicus*. The flora of these strata consists of mixtures of warm and cold-water species. Zone II (SGF 3–10) is characterized by a prominent peak in abundance of the cold-water species *C. pelagicus*. Zone III (SGF 11–18) shows an abrupt decrease of *C. pelagicus* and increase of subtropical to tropical species including *G. oceanica* ($>4\ \mu\text{m}$), *C. leptoporus*, *U. sibogae*, *S. pulchra*, and *C. cristatus*. The change from a cold- to warmer-water flora and increase of the oceanic species took place at SGF 11. Zone IV (SGF 19, 85S30) reveals dominance of *G. oceanica* ($>4\ \mu\text{m}$), *G. caribbeanica* ($>3.5\text{--}4\ \mu\text{m}$), small *Gephyrocapsa* spp. ($<3\ \mu\text{m}$), and *U. hultburtiana* and was barren of *C. pelagicus*.

2. Nannofossil biochronology

The Seoguipo Formation yields Pleistocene marker species such as *G. caribbeanica* ($>3.5\text{--}4\ \mu\text{m}$), *G. oceanica* ($>4\ \mu\text{m}$), *P. lacunosa*, *C. macintyreii*, and *R. asanoi* (Table 2, Figures 2, 3). Among them, the first appearance datum of *G. caribbeanica* (FAD: 1.72 Ma; Sato and Kameo, 1996) has been used for the determination of the Pliocene–Pleistocene boundary. This species is first recognized in sample SGF 5. *G. oceanica* (FAD: 1.65 Ma; Sato and Kameo, 1996), another important indicator for the Pliocene–Pleistocene boundary, occurs in sample SGF 7. *C. macintyreii* with the last occurrence datum (LAD) of 1.60–1.64 Ma (Shipboard Scientific Party, 1996) was encountered in sample SGF 5, although the total occurrence of *C. macintyreii* may not have been properly detected because of its rarity in the samples, and would be expected to occur also in somewhat younger samples. Compiling all these data, the Pliocene–Pleistocene boundary can be drawn between SGF 4 and SGF 5. Consequently, the lowermost part (SGF 1–4) can be presumed to be older than 1.72 Ma and thus belongs to the latest Pliocene.

The sample SGF 11 indicates the lower boundary of Zone CN14a defined by the reoccurrence of *G. oceanica* ($>4\ \mu\text{m}$; 0.94–1.02 Ma; Shipboard Scientific Party, 1996). The presence of *P. lacunosa* (LAD: 0.41 Ma; Sato and Kameo, 1996) indicates that the uppermost part of the formation is older than 0.41 Ma in age. Specimens of *R. asanoi*, with FAD of 1.16 Ma (Sato and Kameo, 1996), are first found in sample SGF 10 at the end of the cooling phase. Therefore, the upper section of sample SGF 10 is younger than 1.16 Ma. This species is last recorded in sample SGF 18, near the

upper boundary of the formation, suggesting that this layer is older than the LAD of *R. asanoi* (0.85 Ma; Sato and Kameo, 1996).


Thus, it is clear that the Seoguipo Formation comprises the latest Pliocene to Early Pleistocene nannofloras. The lower boundary is older than 1.72 Ma, while the upper one is slightly younger than 0.85 Ma. The calcareous nannofossil of this formation is assigned to the *P. lacunosa* Zone (NN19) (Martini, 1971), which corresponds to the combined zones *E. annula* to *E. ovata* (CN13a–CN14a; Okada and Bukry, 1980) of the latest Pliocene to Early Pleistocene.

3. Paleooceanography

The four ecostratigraphic zones established on the basis of the floral composition reflect paleooceanographic conditions (Figure 2). The lowermost section, Zone I is characterized by co-existence of cold (subpolar) and warm (subtropical) water masses. The cooler-water influence increases upward reaching its peak in the middle part of the core (Zone II). It deteriorates toward the upper part (Zone III) until it is totally replaced in Zone IV under the influence of warmer water. This is interpreted as a cooling phase which began in Zone I, culminated in the Zone II, and finally faded out at the upper section (Zone III and IV). In view of the absence of any cold-water current in the present sea around the study area this would seem to be a reflection of the glacial–interglacial cycle rather than due to the local influence of a warm-water current. Comparing the cooling phase with biochronological data, it is certain that a glacial cycle in the study area started shortly before 1.72 Ma (FAD of *G. caribbeanica*) and halted around 0.94–1.02 Ma (reoccurrence of *G. oceanica*).

The distribution of calcareous nannofossils in the Seoguipo Formation also records the position of an oceanographic frontal boundary between warmer water derived from a branch of the Kuroshio Current as it entered the Japan Sea to the north and cooler water introduced into the western portion of the Japan Sea derived from the Liman Current. This oceanographic front oscillated back and forth in response to the ongoing cooling phase. It probably lay near Cheju Island during deposition of Zone I and Zone III, whereas it stood south and north of the island during formations of Zone II and Zone IV, respectively.

Figure 3. All illustrations are light micrographs. Abbreviations: XP=cross-polarized light, TR=transmitted light. Magnification= $\times 2,000$. 1, 2. small *Gephyrocapsa* spp., SGF 12, XP. 3. small *Gephyrocapsa* spp., SGF 11, XP. 4. *Gephyrocapsa caribbeanica*, SGF 5, XP. 5. *Gephyrocapsa caribbeanica*, SGF 10, XP. 6. *Gephyrocapsa oceanica*, SGF 7, XP. 7. *Gephyrocapsa oceanica*, SGF 30, XP. 8, 9. small *Reticulofenestra* spp., SGF 10, XP. 10. small *Reticulofenestra* spp., SGF 11, XP. 11. *Reticulofenestra asanoi*, SGF 15, XP. 12. *Pseudoemiliania lacunosa*, SGF 11, XP. 13. *Pseudoemiliania lacunosa*, SGF 5, XP. 14. *Syracosphaera* sp., SGF 12, XP. 15. *Syracosphaera pulchra*, SGF 18, XP. 16. *Helicosphaera* sp., SGF 12, XP. 17. *Helicosphaera* sp., SGF 17, XP. 18. *Umbilicosphaera sibogae*, SGF 11, XP. 19. *Coccolithus* sp., SGF 16, XP. 20a, b. *Calcidiscus leptoporus*, SGF 11, a: XP., b: TR. 21. *Helicosphaera carteri*, SGF 11, XP. 22. *Helicosphaera carteri*, SGF 12, XP. 23. *Coccolithus* sp., SGF 7, XP. 24. *Coccolithus* sp., SGF 1, XP. 25a, b. *Calcidiscus macintyreii*, SGF 4, a: XP., b: TR. 26. *Calcidiscus macintyreii*, SGF 5, XP. 27. *Calcidiscus leptoporus*, SGF 3, XP. 28. *Coccolithus pelagicus*, SGF 5, XP. 29. *Coccolithus pelagicus*, SGF 7, XP. 30a, b. *Pontosphaera japonica*, SGF 12, a: XP., b: TR. 31. *Pontosphaera japonica*, SGF 7, XP. 32. *Braarudosphaera bigelowii*, SGF 5, XP. 33a, b. *Ceratolithus cristatus*, SGF 4, a: XP., b: TR.



Discussion

1. Stratigraphy

Although the exposure of the Seoguipo Formation above sea level measures about 50 m thick in the Seoguipo area, this formation is developed in the subsurface throughout Cheju Island except for the eastern area of a line connecting Bukchon and Pyoseon. The upheavals of the Seoguipo Formation at Seoguipo City indicate a 150 m uplift of this area at least (cf. Yoon, 1988). However, it is reasonable to suspect that the Seoguipo Formation was formed at a much lower level than where it is at present, since sea level dropped 85–130 m during intense global glacial periods (Muza, 1992).

Quaternary studies on the Seoguipo Formation reported that the outcrop of the Seoguipo Formation in Seoguipo stratigraphically differs from its subsurface strata (Yun *et al.*, 1987; Lee *et al.*, 1987). The above-mentioned researchers considered the age of the Seoguipo Formation to be Middle Pleistocene (0.60–0.41 Ma). They further insisted that subsurface marine sediments recovered from core are Lower Pleistocene (1.2–0.87 Ma) in age (Table 1). For the age determination of sedimentary bodies they applied the paleomagnetic data and radiometric age (K–Ar) of the volcanic rocks that they believed to underlie or overlie two different sedimentary sequences. However, the FAD and LAD of index nannofossils such as *G. caribbeanica* (3.5–4 μm), *G. oceanica* (>4 μm), *C. macintyreii*, *R. asanoi*, *P. lacunosa* and the characteristic nannoflora of the four ecostratigraphic zones recognized in the outcrop section are also observed in 20 cores of Cheju Island. These facts imply that the outcrop strata are to be stratigraphically correlated to subsurface ones. The ecostratigraphic zones and their order of appearance, cold-dominant transitional, cold, warm-dominant transitional, and warm phase from bottom to top can be used as a good stratigraphic tool in Cheju Island. The subdivision of these units largely coincides with that of lithostratigraphic units.

The geologic age and stratigraphic range of the Seoguipo Formation provide a clue in determining the stratigraphic position of comparable sedimentary deposits in Cheju Island and Japan. In previous molluscan studies *Turritella saishuensis*, which occur throughout the Seoguipo Formation, is regarded as an index species for the depositional period of the Seoguipo Formation. However, the lower boundary of this formation is ambiguous due to the absence of stratigraphic marker fossils that may be caused by an influence of cold-water current. Therefore, the FAD of *Turritella saishuensis* also could not be clearly delineated. Some recent paleontological works attempted to define the age of the Seoguipo Formation by paleoceanographic conditions in the Early Pleistocene, when the cold-water mass expanded and

the southward migration of the boreal species proceeded further along the Japan Sea borderland (Lee, 1990; Kang, 1995; Woo *et al.*, 1995). However, there are two different results for the age of first occurrence of *Turritella saishuensis* in the Seoguipo Formation; Pliocene or Pleistocene. The Pliocene to Pleistocene Omma-Manganian fauna in Honshu Island of Japan is correlated with the Seoguipo fauna based on the *Turritella saishuensis* bioseries (Ogasawara, 1981, 1986, 1996; Yoon, 1988). Our result confirms the lower stratigraphic range of the Seoguipo Formation and extension of the FAD of *Turritella saishuensis* to the latest Pliocene. Consequently, the formations correlated to the Seoguipo Formation might be stratigraphically placed between upper Pliocene and Pleistocene (Yoon, 1988).

The paleomagnetic normal polarities of the Seoguipo Formation have been interpreted as the Gauss Normal Polarity Epoch of late Pliocene age (Min *et al.*, 1986) or the Brunhes Normal Polarity Epoch in the middle Pleistocene (Yun *et al.*, 1987). Referring to the reverse polarity of the unconformably overlying Sanbangsan Trachyte, the Seoguipo Formation was placed in the Matuyama Reverse Polarity Epoch of the early Pleistocene (Taneda *et al.*, 1970). However, in a recent paleomagnetic study of the Seoguipo Formation two reversals were recognized in the upper part of the Seoguipo outcrop (Lee, 1998).

2. Biochronology

Previous paleontological studies on the Seoguipo Formation suggested that it was deposited during the Late Pliocene (Yokoyama, 1923; Kim, 1972; Kim, 1984; Yoon, 1988), Pleistocene (Haraguchi, 1931; Lee, 1990; Kang, 1995) or around the Plio-Pleistocene boundary (Paik and Lee, 1984; You *et al.*, 1987).

Our study suggests the age of the upper boundary of the formation is slightly younger than 0.85 Ma based on the presence of *P. lacunosa* (Last Appearance Datum: 0.41 Ma) and LAD (0.85 Ma) of *R. asanoi* (Table 2, Figure 2). It is confirmed by the reoccurrence of *G. oceanica* (>4 μm ; e.g. medium *Gephyrocapsa*) whose acme zone is known to be 0.94–1.02 Ma, in the upper part of the formation. This age of the upper boundary coincides with the results of radiometric K–Ar age of overlying basalt rocks (Table 1; Sanbangsan Trachyte: 0.733 ± 0.056 Ma [Won *et al.*, 1986], 0.87 ± 0.13 Ma [Yun *et al.*, 1987]; Gaksuam Trachyte: 0.893 ± 0.027 Ma [Lee, 1994]). The boundary of the Pleistocene–Pliocene determined by the FAD of *G. caribbeanica* (1.72 Ma) is regarded to lie between Sample SGF 4 and SGF 5. The two samples (SGF 3 and 4) of cold-water environment below Sample SGF 5 are provisionally assigned to the Pliocene, because they may not contain the Pleistocene index taxa for environmental reasons. However, the lowermost part of the formation (Sample SGF1 and 2) definitely belongs to the

Figure 4. All illustrations are Scanning Electron Micrographs. Scale bar equals 2 μm . 1, 2. small *Gephyrocapsa* spp., SGF 11. 3. small *Reticulofenestra* spp., SGF 11. 4. *Pseudoemiliania lacunosa*, SGF 30. 5. *Pseudoemiliania lacunosa*, SGF 11. 6. *Umbilicosaphera sibogae*, SGF 11. 7, 8. *Gephyrocapsa caribbeanica*, SGF 30. 9, 10. *Gephyrocapsa oceanica*, SGF 30. 11. *Coccolithus* sp., SGF 1. 12. *Calcidiscus leptoporus*, SGF 11. 13. *Calcidiscus leptoporus*, SGF 11. 14. *Calcidiscus leptoporus*, SGF 11. 15. *Coccolithus pelagicus*, SGF 5. 16. *Helicosphaera carteri*, SGF 11. 17. *Pontosphaera japonica*, SGF 30. 18. *Syracosphaera pulchra*, SGF 30.

Pliocene in age, since its relatively warmer environment exclude the possibility of climatic effect on distribution of index taxa. This presumption is also supported by normal polarity (Olduvai Normal Polarity Subepoch) specifically obtained from the paleomagnetic analysis of samples SGF 1-SGF 4 (Lee, 1998).

3. Paleoceanography

1). Paleotemperature.—Most paleontological studies reported that the Seoguipo floras and faunas are composed of mixtures of warm- and cold-water elements (Paik and Lee, 1984; You *et al.*, 1987; Yoon, 1988; Lee, 1990; Kang, 1995). According to our results the Seoguipo sequence is characterized by a cooling phase in the lower part and a warm phase in the upper part, and by two transitional phases (Figure 2). Ecostratigraphic Zone I is a first transitional phase in the lowermost part of the formation before beginning of cooling phase. Zone II is characterized by abundant occurrence of *C. pelagicus* is the coldest zone among the eco-zones of the Seoguipo Formation. *C. pelagicus* is presently restricted to the subarctic and transitional waters in the North Atlantic (7°–14°C: McIntyre and Bé, 1967) and North Pacific (6°–14°C: McIntyre *et al.*, 1970). This fact indicates that the paleotemperature of the surface water during deposition of this zone was below 14°C. Zone III is the second transitional phase indicating turnover to the general warming trend. The remarkable bio-event during deposition of the Seoguipo Formation is the sudden emergence of warm-water taxa in this zone. The abundance of *C. pelagicus* abruptly decreases, whereas diversity and abundance of the warm-water elements of *C. leptoporus*, *U. sibogae*, *G. oceanica*, and *S. pulchra* (the first two being oceanic elements) increase substantially. *G. oceanica* is presently distributed within a temperature range of 18°–29°C (McIntyre *et al.*, 1970; Honjo, 1977) or 19°C–31°C (Brand, 1994, p. 44). *Umbilicosphaera sibogae* prefers moderately nutrient-rich, tropical water (Roth and Coulbourn, 1982; Brand, 1994). *S. pulchra* is also a characteristic species in tropical and subtropical environments (Roth and Coulbourn, 1982). Zone IV in the uppermost part of the Seoguipo Formation contains diverse and abundant warm-water taxa. This means that the paleotemperature of the surface water of Zone III and IV was slightly warmer than the average present temperature, 19.38°C (Kwak *et al.*, 1993).

Study of ostracods also revealed similar paleoclimatic patterns to our results in six established biotopes of the formation; cooling phase in the lower part of the formation and warming trend upward (Lee, 1990). Assemblage of biotope 1 and 2 (Eco-Zones I and II: SGF 1-SGF 10) in the lower part of the formation is dominated by cold-water elements such as *Normanicythere sogwipoensis* and *Kotoraicythere paiki* which are completely replaced by the warm-water elements *Cytheropteron uchioi* and *Cytheropteron abnormis* at biotope 3 (basal part of the Eco-Zone III: SGF 11-SGF 12), and in biotopes 4 and 5 (middle part of the Eco-Zone III: SGF 13 – SGF 14) the cold-water elements slightly increase again. The upper part of biotope 6 (upper part of Eco-Zone IV and higher: SGF 15-SGF 19) is characterized by a decrease in the cold-water elements and the occur-

rence of several warm-water elements. The warm-water ostracods found in the Seoguipo Formation are presently distributed in the South China Sea, southern part of the Yellow Sea and Korean South Sea influenced by warm Kuroshio Current (Lee, 1990).

Amano (1994) investigated the surface temperature of the Japan Sea in the Pleistocene through molluscan assemblages from coastal areas of the Japan Sea of Hokkaido to Cheju Island. He estimated the paleotemperature of the Seoguipo Formation as 20°C warmer than the average present temperature, 18.5°C (Marine Environment Map by Japanese Marine Safety Agency).

Woo *et al.*, (1995) analyzed oxygen isotopes from calcareous skeletons of Mollusca. The paleotemperature of his lithologic Unit 4 (SGF 5-SGF 10) from the Seoguipo Formation is calculated to be 13°C. He considered his lithologic Unit 4 the coldest, and the warmest temperature for the entire sequence was 18°C. Paleotemperature of other Units 1, 9, 11, 13 varies from 14 to 17°C. Based on these data, he concluded that the whole sequence of the Seoguipo Formation was deposited during glaciation. His paleoclimatic interpretation coincides with our results for the lower part of the Seoguipo Formation influenced by cold water.

2). Paleoceanographic front.—The site of the Seoguipo Formation is now located in warm surface waters south of an oceanographic front. However, in response to Pleistocene glacial episodes in the last 1.8 m.y. the cold-water front may have migrated southward over Cheju Island as global climate cooled (Momohara, 1994), the sphere of cooler-water influence expanded, and the effect of the warm-water Kuroshio Current diminished (Muza, 1992). Then the Pleistocene nannofossil association in the Seoguipo Formation could record the oscillation or migration of the oceanographic front over Cheju Island as the Japan Sea and the Korea Strait responded to glacial-interglacial cycles. If this is the case, as the oceanographic front migrated southward over Cheju Island owing to global glaciation, cooler surface water would replace warmer water during the deposition of the Seoguipo Formation. This, in turn, may have resulted in a reduced calcareous nannoplankton diversity dominated by cooler-water forms such as *C. pelagicus* as shown in the lower part of the Seoguipo Formation. A lowering of the sea level around the Cheju area in response to the early Pleistocene glaciation is also expected.

Conversely, when the Seoguipo Formation was located south of the oceanographic front in warmer water, assemblage diversity may have been higher, dominated by warmer-water forms such as *G. oceanica*. Since glacial and interglacial cycles have been the primary influence on oceanography of the Japan Sea, Korea Strait, and the vicinity of Cheju Island during the last 1.8 m.y., the changing paleoceanographic conditions of the Seoguipo Formation indicated by the nannofossil assemblages are primarily interpreted as an expression of global climatic changes.

The migration of the oceanographic front is related to the influence of local warm-water currents and global glaciation in the Pleistocene. As a function of these two factors the oceanographic front migrates southward or northward. When global glaciation set about, the cooler-water front was

built first in the Oki Ridge, Ishikawa-Toyama area and migrated southwards to Cheju Island and the Okinawa Trough about 1.8 Ma (Figure 5). It retreated from the Okinawa Trough at 1.60–1.64 Ma (LAD of *C. macintyreii*), from Cheju Island at 0.94–1.02 Ma (reoccurrence of *G. oceanica*), and from the Oki Ridge and the Ishikawa-Toyama area of Japan at 0.85 Ma. The duration of the cooling phase is shortest in the Okinawa Trough and longest in the Japan Sea. Generally invasion of cooler water took place abruptly in comparison to the relatively slow retreating process. Influence of cooler water that migrated southward is well documented in the samples of Oki Ridge, Ishikawa-Toyama area, Niigata area, Seoguipo Formation and Okinawa Trough (Kobayashi, 1990; Muza, 1992; Ogasawara, 1996; Yi *et al.*, *in press*). Considering the geographic location and present-day oceanography only the Niigata area provides an exceptional case, since a cooling phase in this area appeared later and disappeared earlier than expected for its geographic location. The duration of the cooling phase in the Niigata area is also different from that of the neighboring Ishikawa-Toyama area. Therefore, the influence of a warm-water current (*G. inflata* bed No. 2) in this area is considered to be stronger than global glaciation effect in the

Pleistocene.

Regarding the cooling phase and cold-water influence in the Seoguipo Formation that lasted from ca. 1.8 Ma to 0.94–1.02 Ma, it would appear that the Korea Strait never closed during the first Quaternary glaciation. Hence, the cold-water current could expand through the Korea Strait into the Okinawa Trough (Yi *et al.*, *in press*).

3). Paleobathymetry.—The depositional environment of the Seoguipo Formation was interpreted as a littoral zone of shallow sea based on the foraminifera (Kim, 1972), Mollusca (Yoon, 1988) and brachiopods such as *Coptothyris grayi* and *Terebratalia coreanica* that are indicative of a depth of about 50 m (Kim, 1984). Ostracod studies from the Seoguipo Formation enabled a finer subdivision of six biotopes deposited in a shallow sea ranging from bay, through near-shore, to offshore environments (Lee, 1990). An investigation of molluscan fossils suggests the Seoguipo Formation has been formed under the environment of bay to open sea (Kang, 1995).

Among the nannofossil assemblage, neritic species such as *Braarudosphaera bigelowii*, *Gephyrocapsa oceanica*, and *Syracosphaera* spp. occur consistently throughout the formation except for the upper part (Sample SGF 11), where




Figure 5. Correlation of Plio-Pleistocene cooling phases between the Okinawa Trough, Cheju Island, Oki Ridge, and Japan Sea (East Sea) borderland of Japan (Polarity: Cande and Kent, 1995; *: Muza, 1992; **: Ogasawara, 1996).

oceanic species *Calcidiscus leptoporus* and *Umbilicosphaera sibogae* are common. Occurrences of the oceanic taxa in this upper part are accounted for the global warming trend and consequent rising of eustatic sea level. Ostracods from the sample of SGF 11 also indicate that this section was deposited in the outer shelf at depths from 50 m to 100 m.

It is also presumed that the Seoguiipo Formation was uplifted after deposition, because its present altitude is above the presumed sea level in the Early Pleistocene glaciation.

Acknowledgements

We thank E.H. Lee, Korea University, for offering valuable samples. We are also deeply indebted to Iwao Kobayashi (Niigata University), Ritsuo Nomura (Shimane University), Kenshiro Ogasawara (Tsukuba University), and Hisatake Okada (Hokkaido University), for their critical reading of our paper and valuable comments. This study is funded by a grant from the Korea Sciences Engineering Foundation (95-0703-03-01-3). The financial support of the Basic Science Research Program (BSRI-98-9415) of Ministry of Education, Korea is also appreciated.

References

- Ahn, K.S., Lee, H.K. and Lim, H.C., 1995: A petrological and geochemical study of granites in the Cheju Island. *Economic and Environmental Geology*, vol. 28, no. 5, p. 513-518. (In Korean with English Abstract)
- Amano, K., 1994: An attempt to estimate the surface temperature of the Japan Sea in the Early Pleistocene by using a molluscan assemblage. *Paleogeography, Paleoclimatology, Paleoeology*, vol. 108, p. 369-378.
- Brand, L.E., 1994: Physiological ecology of marine coccolithophores. In, Winer, A. and Siesser, W.G. eds., *Coccolithophores*, p. 39-50, Cambridge University Press.
- Cande, S.C. and Kent, D.V., 1995: Revised calibration of the geomagnetic polarity time scale for the Late Cretaceous and Cenozoic. *Journal of Geophysical Research*, vol. 100, p. 6093-6095.
- Honjo, S., 1977: Biogeography and provincialism of living coccolithophorids in the Pacific Ocean. In, Ramsay, A.T.S. ed., *Oceanic micropaleontology*, p. 951-972, London.
- Haraguchi, K., 1931: Geology of Saishu Island. *Bulletin on the Geological Survey of Chosen (Korea)*, vol. 10, pt. 1, p. 1-34, pl. 1-9 (in Japanese); p. 1-12, pl. 1. (in English)
- Inoue, E., 1982: Geological problems on Cretaceous and Tertiary rocks in and around Tsushima-Korea Straits. *United Nations ESCAP, CCOP Technical Bulletin*, vol. 15, p. 85-121.
- Kang, S.S., 1995: Sedimentary facies and paleoenvironment of the Lower Pleistocene Seogwipo Formation, Cheju Island, Korea. *The Quaternary Research*, vol. 34, no. 1, p. 19-38.
- Kim, B.K., 1972: A stratigraphic and paleontologic study of the Seogwipo Formation. *The Memoirs for Professor Chi Moo Son's Sixtieth Birthday*, p. 169-187. (in Korean with English abstract)
- Kim, B.K., 1984: Pliocene brachiopods from the Seogwipo Formation of Jeju Island, Korea. *Journal of the National Academy of Sciences, Republic of Korea, Natural Sciences Series*, vol. 23, p. 167-190, pls. 4.
- Kobayashi, I., 1990: Plio-Early Pleistocene faunal change in the Paleo-Japan Sea, especially in the Niigata Oil Field Area, Central Honshu, Japan. In, Tsuchi, R. ed., *Pacific Neogene events: Their timing nature and inter-relationship*, p. 77-83. Tokyo University Press.
- Kwak, H.S., Byun, S.K., Kim, C., Lee, D.S., Oh, Y.M., Lee, S.H., Je, J.G., Yoo, H.R. and Shin, D.H., 1993: Oceanographic Atlas of Korea Water, Volume 2. South Sea. Korea Ocean Research Development Institute, p. 1-157.
- Lee, D., 1998: Magnetostratigraphy on drill cores and outcrop samples of the Seoguiipo Formation in Cheju Island, Korea. Unpublished Master Thesis, 91 p. Pusan National University. (in Korean with English abstract)
- Lee, D.Y., Yun, S.K., Kim, J.Y. and Kim, Y.J., 1987: Quaternary geology of the Jeju Island. *Korea Institute of Energy and Resources (KIER), Research Report*, KR-87-29, p. 233-278. (in Korean with English abstract)
- Lee, E.H., 1990: Pleistocene Ostracoda from the marine sedimentary strata of the Cheju Island, Korea. Unpublished Ph.D. Thesis, 400 p. Korea University.
- Lee, M.W., 1982a: Geology of Jeju volcanic Island, Korea. *The journal of the Japanese Association of Mineralogists, Petrologists and Economic Geologists*, vol. 77, 55-64. (in Japanese with English abstract)
- Lee, M.W., 1982b: Petrology and geochemistry of Jeju volcanic Island, Korea. *Science Report of the Tohoku University, Sendai, 3rd series*, vol. 15, no. 2, 177-256.
- Lee, M.W., 1994: Development history and geologic structure of Cheju Island. In, *Proceedings of Environmental Preservation and Development of the Groundwater of Cheju Island*, p. 54-74. The Korean Society of Groundwater and Environmental Science and Center for Mineral Resources Research. (in Korean, title translated)
- Martini, E., 1971: Standard Tertiary and Quaternary nannoplankton zonation. In, Farinacci, A. ed., *Proceedings II Planktonic Conference, Roma, 1970*, vol. 2, p. 739-785.
- McIntyre, A. and Be, A.W.H., 1967: Modern Coccolithophoridae of the Atlantic Ocean. I. Placoliths and cyrtoliths. *Deep-Sea Research*, vol. 14, p. 561-597.
- McIntyre, A., Bé, A.W.H. and Roche, M.B., 1970: Modern Pacific coccolithophorida: a paleontological thermometer. *Transactions New York Academy of Sciences*, vol. 32, p. 720-731.
- Min, K.D., Won, J.S. and Hwang, S.Y., 1986: Paleomagnetic study on the volcanic and sedimentary rocks of Jeju Island. *The Journal of the Korean Institute of Mining Geology*, vol. 19, no. 2, p. 153-163. (in Korean with English abstract)
- Momohara, A., 1994: Floral and paleoenvironmental history from the late Pliocene to middle Pleistocene in and around central Japan. *Palaeogeography, Palaeoclimatology, palaeoecology*, vol. 108, p. 281-293.
- Muza, J.P., 1992: Calcareous nannofossil biostratigraphy from the Japan Sea, site 798 and 799: evidence of an oscillating Pleistocene oceanographic frontal boundary. In, Pisciotto, K., Ingle, J.C.Jr., von Breyman, M.T., and Barron, J. et al., *Proceedings of the Ocean Drilling Program, Scientific Results*, vol. 127/128, pt. 1, p. 155-

169. College Station, TX (Ocean Drilling Program).
- Nomura, R., 1984 : Notes on the cassidulinid foraminifera from Jeju Island, Korea. *The memoirs of the Faculty of Education Shimane University (Natural Science)*, vol. 18, p. 21-23.
- Ogasawara, K., 1981 : Paleogeographic significance of the Omma-Maganian fauna of the Japan Sea Borderland. *Saito Ho-on Kai Museum Research Bulletin*, no. 49, p. 17.
- Ogasawara, K., 1986 : Notes on origin and migration of the Omma-Manganian fauna, Japan. *Transactions and Proceedings of the Paleontological Society of Japan*, no. 29, p. 227-244.
- Ogasawara, K., 1996 : Paleobiogeographic significance of the Omma-Manganian fauna. *HGI Report*, no. 5, p. 245-262.
- Okada, H. and Bukry, D., 1980 : Supplementary modification and introduction of code numbers to the low-latitude coccolith biostratigraphic zonation (Bukry, 1973 ; 1975). *Marine Micropaleontology*, vol. 5, no. 3, p. 321-325.
- Paik, K.H. and Lee, E.H., 1984 : A Plio-Pleistocene ostracod assemblage from the Seogwipo Formation, Cheju Island, South Sea of Korea. In, Park, Y.A. et al. eds., *Marine Geology and Physical Processes of the Yellow Sea*, p. 223-234. *Proceedings of US-KOREA Symposium and Workshop on Yellow Sea, 1984, Seoul*.
- Park, B.K., Lee, E.H. and Lee, K.S., 1994 : Stable isotopic compositions of ostracods in the Sogwipo Formation, Korea. *The Journal of Geological Society of Korea*, vol. 30, no. 2, p. 153-158. (in Korean with English abstract)
- Perch-Nielsen, K., 1985 : Mesozoic calcareous nannofossils. In, Bolli, H.M., Saunders, J.B. and Perch-Nielsen, K. eds., *Plankton Stratigraphy*, p. 330-331. Cambridge University Press.
- Rahman, A., 1992 : Calcareous nannofossil biostratigraphy of Leg 127 in the Japan Sea. In, Pisciotto, K.A., Ingle, J.C. Jr., von Breymann, M.T., Barron, J. et al., *Proceedings of the Ocean Drilling Program, Scientific Results*, vol. 127/128, pt. 1, p. 171-186. College Station TX (Ocean Drilling Program).
- Roth, P.H. and Coulbourn, W.T., 1982 : Floral and solution patterns of coccoliths in surface sediments of the North Pacific. *Marine Micropaleontology*, vol. 7, p. 1-52.
- Sameshima, T., Ogawa, T. and Kashima, N., 1988 : Cheju Island (Korea). *5th International Symposium on Vulcanospeleology Excursion Guide Book*, p. 30-36.
- Sato, T. and Kameo, K., 1996 : Pliocene to Quaternary calcareous nannofossil biostratigraphy of the Arctic Ocean, with reference to late Pliocene glaciation. In, Thiedem J., Myhre, A.M., Firth, J.V., Johnson, G.L. and Ruddiman, W.F. eds., *Proceedings of the Ocean Drilling Program, Scientific Results*, vol. 151, p. 39-59. College Station, TX (Ocean Drilling Program).
- Shipboard Scientific Party, 1996 : Explanatory notes. In, Paull, C.K., Matsumoto, R., Wallace, P.J. et al., *Proceedings of the Ocean Drilling Program, Initial Reports*, vol. 164, p. 13-41. College Station, TX (Ocean Drilling Program).
- Tamanyu, S., 1990 : The K-Ar ages and their stratigraphic interpretation of the Cheju Island volcanics, Korea. *Bulletin of the Geological Survey of Japan*, vol. 41, no. 10, p. 527-537.
- Taneda, S., Tsuji, K. and Nakamura, M., 1970 : Geological, petrological and paleomagnetic studies on the volcanic rocks of Cheju Island. *Bulletin of Volcanological Society of Japan*, vol. 2, no. 15, p. 96-108.
- Won, C.K., 1975 : Study of geologic development and the volcanic activity of the Jeju Island. *Journal of Science, Kun Kook University*, vol. 1, p. 7-48. (in Korean with English abstract)
- Won, J.K., Matsuda, J., Nagao, K., Kim, K.H. and Lee, M.W., 1986 : Paleomagnetism and radiometric age of trachytes in Jeju Island, Korea. *The Journal of the Korean Institute of Mining Geology*, vol. 19, no. 1, p. 25-33.
- Woo, K.S., Cheong, D.K. and Park, B.K., 1995 : Paleogeographic investigation from the calcareous skeleton of the Pleistocene Seogwipo Formation, Cheju Island, Korea. *The Journal of the Korean Society of Oceanography*, vol. 30, no. 3, p. 216-226.
- Yi, S., Yun, H. and Choi, D.K., in press : Late Cenozoic nannofossil biostratigraphy of the Okinawa Trough. *Proceedings of the International Conference on "Neogene evolution of Pacific Ocean gateways" Final Meeting of IGCP-355, Kyongju, Korea, 1997*.
- Yokoyama, M., 1923 : On some fossil shells from the Island of Saishu in the Strait of Tsushima. *Journal of the College of Science, Imperial University of Tokyo*, vol. 44, art. 7, p. 1-9, pl. 1.
- Yoon, S., 1988 : The Seogwipo molluscan fauna of Jeju Island, Korea. *Saito Ho-on Kai Special Publication (Professor T. Kotaka Commemorative volume)*, p. 539-545.
- Yoon, S., Koh, G.W. and Kim, I.T., 1995 : Development history of Cheju Island. *Bulletin of Cheju-do Province Government*, vol. 98, p. 148-164. (in Korean)
- You, H.S., Koh, Y.K. and Kim, J.Y., 1987 : Nannofossils from the Seogwipo Formation in Cheju Island, Korea. *Journal of the Paleontological Society of Korea*, vol. 3, no. 2, p. 108-121.
- Yun, S.K., Han, D.S. and Lee, D.Y., 1987 : Quaternary geology in the Southern part of Jeju Island. *Korea Institute of Energy and Resources, Research Report, KR-86-2 (B)-2*, p. 1-64. (in Korean with English abstract)